

Journal of Advances in Biology & Biotechnology

Volume 28, Issue 1, Page 469-475, 2025; Article no.JABB.129808 ISSN: 2394-1081

# Protective Effect of Umbelliferone in Folic Acid-induced Hematological Toxicity in Male C57BL/6

K Divya <sup>a++\*</sup>, Y Ravi Kumar <sup>b#</sup>, B Swathi <sup>c†</sup>, B Anil Kumar <sup>d#</sup>, P Chandrika <sup>a++</sup>, K Ashwini <sup>a++</sup>, G Shirisha <sup>a++</sup>, S Aasritha <sup>a‡</sup>, V Sravathi <sup>a‡</sup>, D Hanuman <sup>a‡</sup>, D Madhuri <sup>c^</sup>, S Soujanya <sup>e†</sup> and B Rajendar <sup>c†</sup>

<sup>a</sup> CVSc, Rajendra Nagar, Hyderabad, India. <sup>b</sup> Department of Veterinary Pathology, CVSc, Korutla, India. <sup>c</sup> Department of Veterinary Pathology, CVSc, Rajendra Nagar, Hyderabad, India. <sup>d</sup> Department of Veterinary Pharmacology & Toxicology, CVSc, Korutla, India. <sup>e</sup> Department of Veterinary Pharmacology, CVSc, Rajendra Nagar, Hyderabad, India.

#### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

#### Article Information

DOI: https://doi.org/10.9734/jabb/2025/v28i11900

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/129808

> Received: 13/11/2024 Accepted: 15/01/2025 Published: 17/01/2025

Original Research Article

++ P.G Scholors;

*Cite as:* Divya, K, Y Ravi Kumar, B Swathi, B Anil Kumar, P Chandrika, K Ashwini, G Shirisha, S Aasritha, V Sravathi, D Hanuman, D Madhuri, S Soujanya, and B Rajendar. 2025. "Protective Effect of Umbelliferone in Folic Acid-Induced Hematological Toxicity in Male C57BL/6". Journal of Advances in Biology & Biotechnology 28 (1):469-75. https://doi.org/10.9734/jabb/2025/v28i11900.

<sup>#</sup> Assistant Professor & Head;

<sup>&</sup>lt;sup>†</sup>Assistant Professor;

<sup>&</sup>lt;sup>‡</sup> PhD Scholars;

<sup>^</sup> Professor & University Head;

<sup>\*</sup>Corresponding author: E-mail: kodavathdivya4@gmail.com;

## ABSTRACT

**Aims:** Folic acid, also referred to as vitamin B9, functions as a catalyst in mono-carbon metabolism, playing a pivotal position in supporting cellular growth and proliferation. Although beneficial, excessive amounts of folic acid can negatively impact hematological health. Herbal remedies are acquiring accelerated popularity in both developed and developing countries, primarily because they are affordable and have fewer complications. Umbelliferone (UMB), known by the name of 7-hydroxycoumarin, is a pharmacologically active compound derived from coumarin. This yellowish-white, crystalline substance found in various plants, exhibiting a Wide array of potency. Using in vivo experimental models, the study sought to explore the beneficial effects of UMB on haematotoxicity caused by folic acid, with a particular emphasis on examining important haematological parameters.

**Frame Work of the Study:** Twenty-four adult C57BL/6 mice were randomly assigned to four groups. Group 1 served as the placebo control. Group 2 was administered folic acid at a dose of 250mg/kg through intra peritoneum on first day to induce toxicity. Group 3 received only UMB at a dose of 60mg/kg body weight through oral route for 14 days. Group 4 was given folic acid on first day and concurrently treated with UMB for 14 days. After 14-day period, hemotological parameres including Total Erythrocyte Count (TEC), Total Leukocyte Count (TLC), Haemoglobin (Hb) concentration, Packed Cell Volume/Haematocrit (PCV/Hct), and Mean Corpuscular Volume (MCV), were meticulously evaluated.

**Results:** Analysis of haematological parameters revealed a significant (p<0.05) decrease in the mean values of TEC, Hb, PCV, MCV, and lymphocytes, along with a marked increase in TLC and neutrophils in group 2 mice compared to other groups. No significant (p<0.05) differences were found between groups 1 and 3. Group 4 showed a mild to moderate improvement in all parameters compared to group 3.

**Conclusion:** This study demonstrates that unambiguously revealed that UMB provides safeguarding counter to haematological toxicity induced by folic acid suggesting its potential as a therapeutic agent.

Keywords: Phytochemicals; one carbon metabolism; oxidative stress; neutrophiles.

# 1. INTRODUCTION

Folates, water-soluble derivatives of vitamin B9, serve as Methyl donors in the Cellular synthesis of thymidylate and purines (Samodelov et al., 2019). Yeast, green vegetables, animal liver and egg volks are sources of folic acid (Pertiwi et al., 2022). Folic acid is essential for rapidly growing tissues, including foetal development, bloodforming organs and epithelial cells (Singh et al., 2024). Due to its key contribution in the synthesis of RNA and DNA components, as well as in DNA methylation, blocking the folate pathway is considered one of the most potent strategies for anti-cancer therapy (Zarou et al., 2021). In spite folic acid supplementation is worthwhile for health, Elevated doses can cause harmful effects on various organs, particularly the kidneys and liver (Al Shawoush et al., 2022). Folic acid is essential for many functions, but excessive intake can mask vitamin B12 deficiency, disrupt methylation, and affect red blood cell production. It may also deplete serum holotranscobalamin, further reducing vitamin B12 availability for critical metabolic processes (Shulpekova et al.,

2021). Serum folate concentrations below 0.005 µg/mL are considered indicative of folate deficiency, while levels below 0.002 µg/mL typically lead to megaloblastic anaemia. Folates generally exhibit low toxicity, with high LD50 values. The mean LD50 for intraperitoneally (i.p.) administered folic acid ranges from 85 to 330 mg/kg body weight (b.wt.) across different mouse strains (Yan, 2021). Phytoconstituents, or phytochemicals, are commonly utilized in traditional medicinal practices and are obtained from plant sources (Basist et al., 2022). On a global scale, numerous phytochemicals and folk remedies, along with their origins, are being studied to verify their effectiveness in alleviating a wide range of afflictions (Chaachouay & Zidane, 2024). Coumarins are natural phenolic widely present in numerous substances medicinal plants (Hassanein et al., 2021). Umbelliferone (UMB) is a coumarin compound recognized for its antioxidant properties and potent anti-inflammatory properties (Liang et al., 2021). Umbelliferone is naturally derived from various plant sources, including coriander (Coriandrum sativum), carrot (Daucus carota),

and garden angelica (*Angelica archangelica*) (Althunibat et al., 2022). The therapeutic profile of umbelliferone is notable for its antidiabetic, anti-inflammatory, anti-nociceptive, anticancer, as well as hepato and nephroprotective attributes (Ali et al., 2021), highlighting its impact in clinical studies and drug discovery. Haematological examination is an essential tool for evaluating the harmful effects of drugs on living organisms. It provides crucial information about metabolites and other components within the body, playing a key role in assessing the animal's physiological, nutritional, and pathological condition (Farag & Alagawany, 2018).

# 2. MATERIALS AND METHODS

# 2.1 Animals

A total of 24 C57BL/6 mice, weighing around 20-25 g, aged 8-12 weeks, were procured from National Institute of Nutrition, Hyderabad. The animals were maintained with an temperature ranging from 22 to 24°C and relative humidity of 50-55% with 12-hour light/dark cycle in propylene cages with each cage containing 3 animals. The mice were allowed a one-week habituation period before the trial commenced. During the study, the animals were given a regular commercial sterilized pellet diet and had accessory to water at leisure. The Institutional Animal Ethics Committee certified the experimental method approval with no (02/28/C.V.Sc., Hyd. IAEC). The mice were equally allocated into four groups, each containing six mice, and the experiment lasted for 14 days. Each group received the assigned treatment according to the schedule outlined below.

# 2.2 Experimental Design

#### List 1. Treatments Details

Group	Treatments
G-I	Placebo control (Normal saline P/O for
	14 days).
G- II	Folic acid (@ 250 mg/kg) I/P single
	dose on the 1 <sup>st</sup> day of experiment.
G-Ⅲ	Umbelliferone per se (@60 mg/kg) P/O
	daily for 14 days.
G-IV	Folic acid (@ 250 mg/kg) I/P single
	dose on the 1 <sup>st</sup> day + Umbelliferone
	(@60 mg/kg) P/O daily for 14 days.

Sample collection and analysis: On 15<sup>th</sup> day the experimental rodents were fasted for 12 hours. Blood was aseptically collected from the retro-orbital plexus of each animal using a capillary tube and transferred into K3-EDTA vacutainers (13 mm x 75 mm, 4 mL, Rapid Diagnostics Pvt. Ltd., Delhi). This ensures the blood is preserved in an anticoagulated state until further analysis. The collected blood samples were utilized for Total Erythrocyte Count (TEC-Millions/µL), Haemoglobin (Hb-a%) Packed concentration. Cell Volume/ Haematocrit (PCV/Hct-percent) and Mean Corpuscular volume (MCV-fL), Total Leukocyte Count (TLC-Thousands/µL), lymphocytes and neutrophils by using automatic whole blood (Huma count. Med Source analvzer Ltd.. Faridabad. Ozone Biomedical Pvt. Harvana) and the results were organized in a systematic table for subsequent statistical analysis.

Statistical analvsis: The data collected were analyzed quantitatively using one-way analysis of variance (ANOVA) with GraphPad Prism 5, version 5.01. (GraphPad Software, California, USA). To discern differences between means, Tukey's test, a multiple comparison procedure, was employed, with the significance level set at p<0.05 (Snedecor & Cochran, 1994).

# 3. RESULTS

Analysis of haemotological parameters revealed a significant decline in TEC. Hb. PCV. MCV and lymphocytes in group II when compared to other groups, however there is considerable increase in TLC and neutrophil count in group II. Statistically, no significant (P<0.05) difference observed between group I and III. A moderate improvement in all these parameters observed in group IV. Any substance that is new to the body is recognized as an antigen, prompting the body to respond and adapt. While umbelliferone generally is considered a safe product, its introduction to the animal results sliaht changes in in parameters when compared to group 1, even though these changes are statistically nonsignificant.



Fig. 1. Ameliorative effect of Umbelliferone on Total Erythrocyte count (TEC)



Fig 3. Ameliorative effect of Umbelliferone on packed cell volume (PCV)



Fig. 5. Ameliorative effect of Umbelliferone on total leukocyte count (TLC)



Fig. 2. Ameliorative effect of Umbelliferone on Haemoglobin concentration



Fig. 4. Ameliorative effect of Umbelliferone on mean cell volume (MCV)



Fig. 6. Ameliorative effect of Umbelliferone on neutrophiles (%)

3.1 Graphical Representation of Effect of UMB on Various Haematological Parameters

472



Fig. 7. Ameliorative effect of Umbelliferone on lymphocytes (%)

### 4. DISCUSSION

Blood is a vital and adaptive component of body that serves as a reflection of the body's overall health, providing insights into both normal physiological processes and the presence of dysfunction disease or (Poletaev, 2018). Haematological assessments play a crucial role in tracking disease progression and evaluating treatment efficacy (Wästerlid et al., 2022). Changes in blood parameters frequently manifest earlier than clinical symptoms, offering an early indication of underlying health conditions. In the present study, a notable (P<0.05) decreased mean values of TEC, Hb, PCV, MCV, Lymphocytes and increased mean values of TLC and neutrophiles were observed in group II when compared with group I. These results align with and are supported by the findings of earlier studies (Vagdevi et al., 2024). Excessive folic disrupts the one-carbon metabolism acid pathway, causing imbalance in purine and pyrimidine synthesis, which subsequently interferes with normal haematopoiesis. Elevated folic acid levels can interfere with Regular metabolic activities, possibly leading to mild inflammation or an increase in oxidative stress (Nikolic et al., 2020). This disruption can impact various biological mechanisms, leading to the activation of the immune system which may trigger activation of inflammatory pathways. This inflammatory condition can trigger the release of cytokines, which enhance the production and mobilization of neutrophils from the bone marrow (Mitroulis et al., 2020). The significant (p<0.05) increase in the mean values of TLC and neutrophils may be attributed to the production of reactive oxygen species (ROS), reflecting a heightened inflammatory response that activates the animal's defense mechanism, resulting in the

influx of inflammatory cells (Sravathi et al., 2022). It is indicative of leukocytosis, which is attributed to enhanced immune function to protect the animals from folic acid-induced toxicity (Hao et al., 2022). In comparison, Group IV mice exhibited a significant rise in TEC, Hb, PCV, MCV, and lymphocyte levels, along with a decrease in the mean values of TLC and neutrophils compared to Group 11. This improvement likely attributed is to the compound's antioxidant and anti-inflammatory properties (Icoglu Aksakal et al., 2021). The antioxidant property of UMB is through upregulation of nrf2 pathway. The antiinflammatory effects of umbelliferone are primarily mediated through the inhibition of the NF-kB pathway (Hassanein et al., 2021).

### 5. CONCLUSION

In conclusion, folic acid leads to a marked decrease in TEC, Hb, PCV, MCV, and lymphocyte levels, along with reductions in TLC and neutrophil counts. In Group IV mice, supplementation with umbelliferone led to a significant (P < 0.05) increase in TEC, Hb, PCV, and MCV, along with a significant (P < 0.05) decrease in TLC levels when compared to Group II mice. The current study demonstrated the protective effect of umbelliferone in mitigating toxicity caused by folic acid. It can be concluded that umbelliferone alleviates cytokine storms and oxidative stress by suppressing the activation of the NF-kB signalling pathway while enhancing the regulation of the Nrf-2 pathway. To deepen our understanding and mitigate haematotoxicity. as well as improve therapeutic outcomes, further in-depth studies on the mechanisms of folic acidinduced haematotoxicity are crucial. Understanding the mechanism of folic acid

toxicity is an important area of research, especially in relation to its metabolism and potential effects at high doses.

#### DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative Al technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

# ACKNOWLEDGEMENTS

The authors are thankful to the College of Veterinary Science, Rajendra nagar, Hyderabad. for according permission to carry out the research work.

# **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

# REFERENCES

- Al Shawoush, A. M., Said, R. S., Hassan, F. E., Ali, S. B., Mohamed, A. S., Elbatran, M. M. (2022). Therapeutic effect of *Nigella sativa* extract on folic acid-induced acute hepatorenal injury: Influences and underlying mechanisms. *Current Topics in Pharmacology, 26*, 49-55. https://doi.org/10.2174/1568026626262203 010000
- Ali, F. E., Hassanein, E. H., El-Bahrawy, A. H., Omar, Z. M., Rashwan, E. K., Abdel-Wahab, B. A., Abd-Elhamid, T. H. (2021). Nephroprotective effect of umbelliferone against cisplatin-induced kidney damage is mediated by regulation of NRF2, cytoglobin, SIRT1/FOXO-3, and NF-kB-p65 signaling pathways. *Journal of Biochemical and Molecular Toxicology*, 35(5), e22738.

https://doi.org/10.1002/jbt.22738

Allam, M. A., Khowailed, A. A., Elattar, S., & Mahmoud, A. M. (2022). Umbelliferone ameliorates oxidative stress and testicular injury, improves steroidogenesis and upregulates peroxisome proliferatoractivated receptor gamma in type 2 diabetic rats. *Journal of Pharmacy and Pharmacology*, *74*(4), 573-584.

https://doi.org/10.1002/jphp.13625

Althunibat, O. Y., Abduh, M. S., Abukhalil, M. H.,
Aladaileh, S. H., Hanieh, H., & Mahmoud,
A. M. (2022). Umbelliferone prevents
isoproterenol-induced myocardial injury by

upregulating Nrf2/HO-1 signaling, and attenuating oxidative stress, inflammation, and cell death in rats. *Biomedicine & Pharmacotherapy*, *149*, 112900. https://doi.org/10.1016/j.biopha.2022.1129 00

Basist, P., Parveen, B., Zahiruddin, S., Gautam, G., Parveen, R., Khan, M. A., Krishnan, A., Shahid, M., & Ahmad, S. (2022). Potential nephroprotective phytochemicals: Mechanism and future. *Environmental Toxicology and Pharmacology, 283*, 114743.

https://doi.org/10.1016/j.etap.2022.114743

- Chaachouay, N., & Zidane, L. (2024). Plantderived natural products: A source for drug discovery and development. *Drugs and Drug Candidates, 3*(1), 184-207. https://doi.org/10.1007/s12325-023-02347y
- Farag, M. R., & Alagawany, M. (2018). Erythrocytes as a biological model for screening of xenobiotics toxicity. *Chemico-Biological Interactions*, 279, 73-83. https://doi.org/10.1016/j.cbi.2017.11.013
- Hao, Y., Romano, J. D., & Moore, J. H. (2022).
  Knowledge-guided deep learning models of drug toxicity improve interpretation. *Patterns, 3*(9).
  https://doi.org/10.1016/j.patter.2022.10046
- Hassanein, E. H., Ali, F. E., Kozman, M. R., & Abd El-Ghafar, O. A. (2021). Umbelliferone attenuates gentamicin-induced renal toxicity by suppression of TLR-4/NF-κBp65/NLRP-3 and JAK1/STAT-3 signaling pathways. *Environmental Science and Pollution Research, 28*, 11558-71. https://doi.org/10.1007/s11356-020-12075-5
- Icoglu Aksakal, F., Koc, K., Geyikoglu, F., & Karakaya, S. (2021). Ameliorative effect of umbelliferone in remote organ injury induced by renal ischemia-reperfusion in rats. *Journal of Food Biochemistry*, 45(2), e13628. https://doi.org/10.1111/jfbc.13628
- Liang, S., Chen, Z., Li, H., Cang, Z., Yin, K., Wu, M., & Luo, S. (2021). Neuroprotective effect of umbelliferone against cerebral ischemia/reperfusion induced neurological deficits: In-vivo and in-silico studies. *Journal of Biomolecular Structure and Dynamics, 39*(13), 4715-25. https://doi.org/10.1080/07391102.2021.188 4054
- Mitroulis, I., Kalafati, L., Bornhäuser, M., Hajishengallis, G., & Chavakis, T. (2020).

Regulation of the bone marrow niche by inflammation. *Frontiers in Immunology, 11*, 1540.

https://doi.org/10.3389/fimmu.2020.01540

- Nikolic, T., Petrovic, D., Matic, S., Turnic, T. N., Jeremic, J., Radonjic, K., Srejovic, I., Zivkovic, V., Bolevich, S., & Jakovljevic, V. (2020). The influence of folic acid induced acute kidney injury on cardiac function and redox status in Naunynrats. Schmiedeberg's Archives of Pharmacology, 393, 99-109. https://doi.org/10.1007/s00210-020-01967-3
- Pertiwi, H., Nur Mahendra, M. Y., & Kamaludeen, J. (2022). Folic acid: Sources, chemistry, absorption, metabolism, beneficial effects on poultry performance and health. *Veterinary Medicine International, 2022*(1), 2163756.

https://doi.org/10.1155/2022/2163756

- Poletaev, A. (2018). Composition of the blood and reflection of the health state of human body. *Biomedical and Pharmacology Journal*, 11(4), 1797-800.
  - https://doi.org/10.22329/bpj.v11i4.6418
- Samodelov, S. L., Gai, Z., Kullak-Ublick, G. A., & Visentin, M. (2019). Renal reabsorption of folates: Pharmacological and toxicological snapshots. *Nutrients, 11*(10), 2353. https://doi.org/10.3390/nu11102353
- Shulpekova, Y., Nechaev, V., Kardasheva, S., Sedova, A., Kurbatova, A., Bueverova, E., Kopylov, A., Malsagova, K., Dlamini, J. C., & Ivashkin, V. (2021). The concept of folic acid in health and disease. *Molecules*, *26*(12), 3731. https://doi.org/10.3390/molecules2612373
- Singh, A. P., Maurya, N. K., Saxena, R., & Saxena, S. (2024). An overview of red blood cell properties and functions. *Journal*

of International Research in Medical and Pharmaceutical Sciences, 19(2), 14-23. https://doi.org/10.47607/jir-19-2-14

- Snedecor, G. W., & Cochran, G. (1994). Statistical methods (8th ed.). Iowa State University Press.
- Sravathi, V., Jeevanalatha, M., Ravikumar, Y., Gopalareddy, T., Chandravathi, B. S., & Lakshman, M. (2022). 5-Fluorouracil induced hematological alterations in Wistar rats and its amelioration with naringenin. *Pharma Innovation Journal, SP-11*(1), 1250-1254.
- Vagdevi, T., Ravikumar, Y., Chandravathi, T., Kumar, B. A., Haripriya, B., Sagar, S., Aashritha, S., Sravathi, V., Hanuman, D., Preethi, B., & Swathi, B. (2024). Folic acid induced Haematotoxicity in mice and its amelioration with Ferulic acid. *International Journal of Advanced Biochemistry Research, 8*(1), 828-831.

https://doi.org/10.58262/ijabr.2024.1019

- Wästerlid, T., Cavelier, L., Haferlach, C., Konopleva, M., Fröhling, S., Östling, P., Bullinger, L., Fioretos, T., & Smedby, K. E. (2022). Application of precision medicine in clinical routine in haematology challenges and opportunities. *Journal of Internal Medicine, 292*(2), 243-61. https://doi.org/10.1111/joim.13435
- Yan, L. J. (2021). Folic acid-induced animal model of kidney disease. *Animal Models and Experimental Medicine, 4*(4), 329-42. https://doi.org/10.1002/ame2.12112
- Zarou, M. M., Vazquez, A., & Vignir Helgason, G. (2021). Folate metabolism: A re-emerging therapeutic target in haematological cancers. *Leukemia*, *35*(6), 1539-51. https://doi.org/10.1038/s41375-021-01061w

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/129808