



Response of Levels of Nitrogen and Row Spacing on Growth and Leaf Yield of Palak (*Beta vulgaris* var. *bengalensis*) var. Arka Anupama

**M. J. Patel ^{a*}, B. H. Panchal ^a, A. S. Thounaojam ^b,
K. P. Zala ^c and V. K. Baria ^a**

^a Department of Horticulture, B. A. College of Agriculture, Anand Agricultural University, Anand-388 110, Gujarat, India.

^b Medicinal and Aromatic Plants Research Station, Anand Agricultural University, Anand – 388 110, Gujarat, India.

^c Department of vegetable science, College of horticulture, Sardarkrushinagar Dantiwada Agricultural University, Sardarkrushinagar- 385506, Gujarat, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/ijpss/2025/v37i15254>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/129134>

Original Research Article

Received: 28/10/2024

Accepted: 30/12/2024

Published: 11/01/2025

ABSTRACT

A field experiment was conducted at Horticultural Research Farm, Department of Horticulture, B. A. College of Agriculture, Anand Agricultural University, Anand, with a view to study the response of levels of nitrogen and row spacing on growth and leaf yield of palak (*Beta vulgaris* var. *bengalensis*)

*Corresponding author: E-mail: mahimapatel644@gmail.com;

var. Arka Anupama during Rabi season of the year 2019-20 with factorial randomized block design (FRBD) comprising of three levels of nitrogen viz., 60 kg Nha⁻¹ (N₁), 80 kg Nha⁻¹ (N₂) and 100 kg Nha⁻¹ (N₃) and three row spacing viz., 20 cm (S₁), 30 cm (S₂) and 40 cm (S₃) along with three replications. The study was revealed that the application of Nitrogen 100 kgha⁻¹ (N₃) gave maximum shoot length (29.24 cm), maximum number of leaves (13.37), maximum leaf area (91.02 cm²), chlorophyll content SPAD value (37.10), ascorbic acid content (76.47 mg) at first cutting, maximum fresh leaf weight i.e. 57.88, 52.77, 48.23 and 38.33 g and maximum dry leaf weight i.e. 6.50, 6.24, 5.81 and 4.92 g at 1st, 2nd, 3rd and 4th cutting, respectively and maximum leaf yield per plot (20.90 kg) and per hectare (37.32 t) while in case of row spacing, S₃ (40 cm) gave maximum number of leaves (13.44), maximum leaf area (89.69 cm²) at first cutting, maximum fresh leaf weight i.e. 56.63, 51.92, 46.30 and 36.67 g and maximum dry leaf weight i.e. 6.41, 6.20, 5.82 and 4.75 g at 1st, 2nd, 3rd and 4th cutting respectively while, maximum number of plants/sq.m at 10 DAS (47.05), shoot length (28.98 cm), green leaf yield per plot (20.68 kg) and per hectare (36.93 t) observed in treatment S₁ (20 cm).

Keywords: Nitrogen; row spacing; leaf yield; Spinach.

1. INTRODUCTION

Palak (*Beta vulgaris var. bengalensis*) is one of the most important leafy vegetables of tropical and sub-tropical regions. It is also known by other name like leaf beet, beet palak and spinach beet. It is closely related to beetroot (*Beta vulgaris*) and Swiss chard (*Beta vulgaris var. cicla*) and good substitute for spinach. Palak has a nutritional as well as medicinal value. Green leafy vegetables occupy an important among the food crops as these provide adequate amounts of many vitamins and minerals for humans. Amongst all the vegetables, the leafy vegetables have high protective food value. They are rich in mineral and hence called as 'mines of minerals'. Vitamin A and C are present in abundant quantities. Beside this, soft fibrous matter is specially providing necessary roughage in diet. It is widely grown leafy vegetable. The leaves are used in inflammation, paralysis, headache and remedy for liver disease. Besides its medicinal property it also acts as mild laxative. Palak neutralize the acidity produced during digestion of fatty substance and help to prevent constipation. Indian spinach is used as fresh vegetables for cooking and also in salad form. It contains moisture 86.49 %, fiber 0.7 g, protein 3.4 g, minerals 2.2 g, carbohydrates 6.5 g, riboflavin 0.5 g, calcium 380 mg, iron 16.2 mg, thiamin 0.26 g, Vitamin A 9770 IU, Vitamin C 70 mg/ 100g of edible portion (Vishnu, 2014).

The several factors like judicious fertilization and use of optimum amount of quality seed play an important role in crop production. Spinach beet being a green leafy vegetable may need heavy nitrogen fertilization but its exact requirement not known and hence it is to be examined. Nitrogen

in general increases the vegetative growth but depresses the root growth. Excess of nitrogen strongly stimulates the plant to synthesize proteins and to develop fresh vegetative tissues and the bulk of the carbohydrates is used for the formation of amino acid and proteins and the construction of strengthening tissues, in consequence to become a spongy, tender and dark green.

Nitrogen is the essential for the vegetative growth of the plant resulting in higher green leaf yield. An ample supply of nitrogen not only helps in the production of succulent leafy matter in all leafy vegetable but also in their seed production. The yield of palak depends on vegetative growth and it may express in terms of number of leaves per plant, size of leaf, leaf area and plant height. For obtaining more vegetative growth cutting of crop is important due to cutting of crop side shoots are arises which increases the number of leaves per plant and ultimately increased the yield which demands higher amount of nitrogen.

Spacing between rows of plants is another factor that affects the growth and yield of the crop. Evapotranspiration and weed infestation were found high in the crop grown with wider spacing and hence it is necessary to grow the crop at optimum spacing (Rahman & Talukdar 1986). In palak not only planting geometry but also nutrient management practices play a vital role in boosting up its productivity. Optimum row spacing facilitates better utilization of nutrient resources supplied through inorganic fertilizers so keep in view study was carried out to determine the response of levels of nitrogen and row spacing and their combination on growth and yield of palak.

2. MATERIALS AND METHODS

Present experiment was conducted in open field condition at Horticultural Research Farm, Department of Horticulture. B. A. College of Agriculture, A. A. U., Anand, Gujarat during *Rabi* season of the year 2019-20 in which efforts were made to investigate the growth and leaf yield of palak with different levels of nitrogen and row spacing. The experiment was layout in Factorial Randomized Block Design (FRBD) with three replications comprising three levels of nitrogen viz., 60 kg Nha⁻¹ (N₁), 80 kg Nha⁻¹ (N₂) and 100 kg Nha⁻¹ (N₃) and three row spacing viz., 20 cm (S₁), 30 cm (S₂) and 40 cm (S₃) comprising total nine treatment combinations were taken for experiment. Experiment site was well prepared, cultural practices includes thinning, weeding, irrigation and manure application were followed for the healthy growth of crop.

The field was ploughed thoroughly and then rotavated until the soil was friable and free from perennial weeds after that well-rotten farm yard manure was incorporated into the soil at the time of final ploughing. Half dose of nitrogen as per treatment was applied in form of urea along with full dose of phosphorus and potassium fertilizer as basal dose at time of field preparation and remaining half dose of nitrogen was applied after each cutting or harvesting of the leaves. Seeds were sown at 6-7 cm depth at a distance of 10 cm within row by hand and covered with fine soil. Seeds were sown at third week of October. Green leaves of palak were harvested at 40, 55, 70 and 85 days after sowing at 5 cm above ground level. The palak variety Arka Anupama was taken for present investigation. Ten plants per treatment plot were observed and recorded for growth and yield characteristics. Palak growth and yield parameters taken at different stages of harvesting like number of plants per sq.m is measured at ten days after sowing, while shoot length, number of leaves per plant and leaf area were measured at first cutting of leaves and fresh and dry weight of leaves were measured at each cutting. The fresh weight of leaves taken using electrical balance for each treatment separately. For the dry weight, leaves was oven dried at 60 °C for 24 hrs. The experiment data were analyzed using "Analysis of Variance Techniques" in Factorial Randomized Block Design (FRBD). Data on growth and yield parameters were collected and discussed below.

3. RESULTS AND DISCUSSION

3.1 Growth Parameters

3.1.1 Number of plants/ m²

Number of plants/ m² at 10 days after sowing was found non-significant effect with respect to different levels of nitrogen. Among the row spacing treatment S₁ (20 cm) found significant. This might be due to closer spacing increase the number of plants. The results are in conformity with the findings of Snider et al. (2012) in sorghum.

3.1.2 Shoot length (cm)

Shoot length was significantly affected due to various levels of nitrogen. Maximum shoot length (29.24 cm) was observed in treatment N₃ (100 kg Nha⁻¹). This might be due to the fact that, nitrogen which is vital nutrient compound and a major constituent of chlorophyll, chief constituent of protein, essential for the formation of protoplasm which help in stimulating the cell division and cell elongation which leads increased plant height and growth. The results are closely relevant to the findings of Sharma et al. (2016) in coriander. While among the row spacing maximum shoot length (28.98 cm) was observed in treatment S₁ (20 cm), which was at par with treatment S₂ (27 cm). This might be due to more plant population per unit area which resulted in more competition for sunlight leading to more up right growth in search of sunlight. The results are in conformity with the findings of Zibelo et al. (2016) in okra.

3.1.3 Number of leaves/plant

Number of leaves/plant at first cutting was significantly affected by the different levels of nitrogen. Maximum number of leaves (13.37) observed in treatment N₃ (100 kg Nha⁻¹), which was at par with treatment N₂ (13.05) it might be due to higher level of nitrogen imparts the vigorous vegetative growth which resulting in increased number of leaves per plant. Similar results are also reported by Ali et al. (2016) in spinach and Anupama et al. (2017) in fenugreek. Among the row spacing maximum number of leaves (13.44) observed in treatment S₃ (40 cm), which was at par with treatment S₂ (12.82) it might be due to plants get sufficient space for vegetative growth, faced no competition for the sunlight which resulted in more vegetative growth which expressed in production of more number of leaves per plant. Similar, results were also reported by Patel et al. (2011) in amaranthus.

Table 1. Effect of nitrogen and row spacing on number of plants/ m² at 10 DAS, shoot length, number of leaves per plant, leaf area, green leaf yield kg/plot and green leaf yield t/ha of palak var. Arka Anupama

Treatments	Number of plants/ m ²	Shoot length (cm)	Number of leaves per plant	Leaf area (cm ²)	Green leaf yield kg/plot	Green leaf yield t/ha
Factor A : Nitrogen						
N ₁ : 60 kg/ha	34.76	25.79	11.55	74.01	17.91	31.98
N ₂ : 80kg/ha	34.27	27.01	13.05	82.56	18.46	32.97
N ₃ : 100 kg/ha	33.77	29.24	13.37	91.02	20.90	37.32
S. Em.±	1.14	0.73	0.40	3.47	0.64	1.14
C. D. at 5 %	NS	2.19	1.21	10.42	1.92	3.43
Factor B : Row spacing						
S ₁ : 20 cm	47.05	28.98	11.70	75.57	20.68	36.93
S ₂ : 30 cm	31.76	27.00	12.82	82.33	18.69	33.38
S ₃ : 40 cm	23.98	26.06	13.44	89.69	17.90	31.96
S. Em.±	1.14	0.73	0.40	3.47	0.64	1.146
C. D. at 5 %	3.43	2.19	1.21	10.42	1.92	3.436
Interaction						
N × S	NS	NS	NS	NS	NS	NS
C.V. %	10.04	8.03	9.63	12.63	10.09	10.09

Table 2. Effect of nitrogen and row spacing on fresh weight of leaves at 1st, 2nd, 3rd and 4th cutting of palak var. Arka Anupama

Treatments	Fresh leaf weight (g)			
	At cutting	1 st At 2 nd cutting	At 3 rd Cutting	At 4 th Cutting
Factor A : Nitrogen				
N ₁ : 60 kg/ha	51.35	44.22	38.44	31.89
N ₂ : 80kg/ha	54.89	47.61	42.93	33.34
N ₃ : 100 kg/ha	57.88	52.77	48.23	38.33
S. Em.±	1.33	1.74	2.02	1.35
C. D. at 5 %	3.99	5.23	6.07	4.07
Factor B : Row spacing				
S ₁ : 20 cm	51.66	45.02	37.49	31.18
S ₂ : 30 cm	55.84	47.66	45.82	35.71
S ₃ : 40 cm	56.63	51.92	46.30	36.67
S. Em.±	1.33	1.74	2.02	1.35
C. D. at 5 %	3.99	5.23	6.07	4.07
Interaction				
N × S	NS	NS	NS	NS
C. V. %	7.30	10.86	14.06	11.80

3.1.4 Leaf area (cm²)

As regards to leaf area (cm²) per plant at first harvest nitrogen 100 kg/ha⁻¹ (N₃) recorded significantly highest leaf area (91.02 cm²) of palak followed by N₂ (82.56 cm²) it might be due to the quick availability of nitrogen to the plants, nitrogen imparts the vigorous vegetative growth which leads to the cell division and cell enlargement. Similar, results were also reported

by Singh et al. (2015) in palak and Elsayed and Abdelraouf (2008) in spinach. With regards to leaf area at first cutting showed significantly, maximum leaf area (89.7 cm²) in treatment S₃ (40 cm), which was at par with treatment S₂ (82.33 cm²) that might be due to more favourable condition with better uptake of nutrients, water and efficient utilization of sunlight. Similar, results were also recorded by Patel et al. (2011) in amaranthus.

Table 3. Effect of nitrogen and row spacing on dry weight of leaves at 1st, 2nd, 3rd and 4th cutting of palak var. Arka Anupama

Treatments	Dry leaf weight (g)			
	At 1 st cutting	At 2 nd Cutting	At 3 rd Cutting	At 4 th Cutting
Factor A : Nitrogen				
N ₁ : 60 kg/ha	5.20	5.09	4.61	4.01
N ₂ : 80kg/ha	5.90	5.80	5.48	4.38
N ₃ : 100 kg/ha	6.50	6.24	5.82	4.92
S. Em.±	0.26	0.23	0.24	0.19
C. D. at 5 %	0.79	0.69	0.74	0.59
Factor B : Row spacing				
S ₁ : 20 cm	5.40	5.22	4.83	4.01
S ₂ : 30 cm	5.79	5.71	5.26	4.54
S ₃ : 40 cm	6.41	6.20	5.81	4.75
S. Em.±	0.26	0.23	0.24	0.19
C. D. at 5 %	0.79	0.69	0.74	0.59
Interaction				
N × S	NS	NS	NS	NS
C. V. %	13.60	10.86	14.06	13.33

3.1.5 Fresh leaf weight (g)

With respect to fresh leaf weight at 1st, 2nd, 3rd and 4th cutting nitrogen 100 kg ha⁻¹ (N₃) found significantly maximum fresh leaf weight i.e. 57.88, 52.77, 48.23 and 38.33 g respectively which was at par with N₂ (80 kg ha⁻¹) at 1st, 2nd, 3rd and 4th cutting that might be due to the fact that nitrogen is the vital component of protein which promotes cell division and as the level of nitrogen increases, it promotes the vigorous cell division and accumulation of more plant metabolites. Similar, trends in results were also observed by et al. (2016) in coriander and Anupama et al. (2017) in fenugreek. With regards to row spacing fresh leaf weight at 1st, 2nd, 3rd and 4th cutting treatment S₃ (40 cm) recorded significantly maximum fresh leaf weight i.e. 56.63, 51.92, 46.30 and 35.71 g respectively which was at par with S₂ (30 cm) at 1st, 2nd, 3rd and 4th cutting that might be due to the fact that plant receive enough light and nutrients which leads to attain maximum fresh weight of leaves. Similar, results were also reported by Patel et al. (2011) in amaranthus.

3.1.6 Dry leaf weight (g)

With respect to dry leaf weight at 1st, 2nd, 3rd and 4th cutting nitrogen 100 kg ha⁻¹ (N₃) found significantly maximum dry leaf weight i.e. 6.50, 6.24, 5.81 and 4.92 g respectively which was at par with N₂ (80 kg ha⁻¹) at 1st, 2nd, 3rd and 4th cutting that might be due to higher level of nitrogen helps in vigorous vegetative growth and

accumulation of more photosynthetic product in the plants. On drying plants with healthy and vigorous growth resulted in maximum dry matter than that of plants with lower levels of nitrogen. Similar, results were also reported by Biemond (2004) and Pujari (2017) in spinach. With regards to row spacing dry leaf weight at 1st, 2nd, 3rd and 4th cutting treatment S₃ (40 cm) recorded significantly maximum dry leaf weight i.e. 6.41, 6.20, 5.82 and 4.75 g respectively which was at par with S₂ (30 cm) at 1st, 2nd, 3rd and 4th cutting that might be due to plants get sufficient space for vegetative growth, faced no competition for the sunlight which resulted in more vegetative growth which expressed in production of more number of leaves per plant and thereby increase the fresh leaf weight and upon drying increase the dry weight of leaves. These results are closed to finding of Patel et al. (2011) in amaranthus.

3.2 Yield Parameters

3.2.1 Green leaf yield (kg/plot)

Nitrogen 100 kg N ha⁻¹ (N₃) recorded significantly highest green leaf yield per plot (20.90 kg) that might be due to nitrogen play a role of stimulant in metabolic activity which leads to the cell division and cell enlargement and probably increased the number of leaves, leaf area and fresh weight of leaves which finally lead to increased green leaf yield per plot and per hectare. The result was in confirmly with Anupama et al. (2017) in fenugreek, Datta et al.

(2008) in coriander and Elsayed & Abdelraouf (2008) in spinach. With regards to row spacing highest green leaf yield per plot (20.68 kg) was observed in treatment S₁ (20 cm) that to be fact that in close spacing there was more number of plants per unit area of land. As the plants per unit area were increased the green yield per plot increased significantly. The results are in accordance with finding of Essilfie et al. (2017) in chilli and Assefa et al. (2015) in tomato.

3.2.2 Green leaf yield (t/ha)

Nitrogen 100 kg Nha⁻¹ (N₃) recorded significantly highest green leaf yield per hectare (37.32 t/ha) With regards to row spacing highest green leaf yield per hectare (36.93 t/ha) was observed in treatment S₁ (20 cm).

4. CONCLUSION

The several factors like judicious fertilization and use of optimum amount of quality seed play an important role in crop production. Spinach beet being a green leafy vegetable may need heavy nitrogen fertilization but its exact requirement not known and hence it is to be examined. Based on the results, it was concluded that the application of 100 kg N ha⁻¹ combined with a row spacing of 20 cm was most effective for achieving a higher green leaf yield in palak.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Ali, S., Zeb, B., & Khan, T. H. (2016). Impact of different levels of nitrogen on the yield of spinach. *International Journal of Advanced Research and Review*, 1(12), 80-87.
- Anupama, G., Hegde, L. N., Hegde, N. K., Devappa, V., Mastiholi, A. B., & Sandhyarani, N. (2017). Effect of nitrogen and spacing levels on growth and yield parameters of *Kasuri methi* (*Trigonella corniculata* L.) var. Pusa Kasuri. *International Journal of Current Microbiology and Applied Sciences*, 6(9), 1464-1469.
- Assefa, W., Tesfaye, B., & Dessalegn, L. (2015). Influence of inter and intra-rows spacing on yield and yield components of tomato cultivars. *Ethiopia Journal of Agriculture Science*, 25, 71-81.
- Biemond, H. (2004). Effect of nitrogen on development and growth of the leaves of vegetables. Appearance and expansion growth of leaves of spinach. *Pakistan Journal of Biological Science*, 7(1), 82-94.
- Datta, S., Alam, K., & Chatterjee, R. (2008). Effect of different levels of nitrogen and leaf cutting on growth, leaf and seed yield of coriander. *Indian Journal of Horticulture*, 65(2), 201-203.
- Elsayed, A., & Abdelraouf, A. (2016). The effects of nitrogen fertilization on yield and quality of spinach grown in high tunnels. *Alexandria Science Exchange Journal*, 37(3), 488-496.
- Essilfie, M. E., Dapaah, H. K., Boateng, E., & Damoah, R. J. (2017). Age of transplant and row spacing effects on growth, yield and yield components of chilli pepper (*Capsicum annuum* L.). *International Journal of Environment, Agriculture and Biotechnology*, 2(5), 2406-2418.
- Patel, S. P. A., Alagundagi, S. C., Mansur, C. P., Kubsad, V. S., Hosamani, S. V., & Megeri, S. N. (2011). Effect of row spacing and seed rate on growth, fodder productivity and economics of amaranth genotypes. *Karnataka Journal of Agriculture Science*, 24(5), 651-653.
- Pujari, K. (2017). Effect of nitrogen and phosphorous levels on performance of Indian spinach under different growing conditions. [M.Sc. thesis, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola].
- Rahman, A. K. M. M., & Talukdar, M. P. (1986). Influence of date of planting and plant spacing on the growth and yield of garlic. *Bangladesh Journal of Agriculture*, 11, 19-26.
- Sharma, A., Naruka, I. S., & Shakhawat, R. P. S. (2016). Effect of row spacing and nitrogen on growth and yield of coriander (*Coriandrum sativum* L.). *Journal of Krishi Vigyan*, 5(1), 49-53.
- Snider, J. L., Raper, R. L., & Schwab, E. B. (2012). The effect of row spacing and seeding rate on biomass production and plant stand characteristics of non-irrigated photoperiod-sensitive sorghum (*Sorghum*

- bicolor* L. Moench). *Industrial Crops and Products*, 37, 527-535.
- Vishnu, S. (2014). *Vegetable Science and Technology in India*. Kalyani Publishers.
- Zibelo, H., Wtsadik, K., & Sharma, J. J. (2016). Effect of inter and intra row spacing on growth and yield of okra (*Abelmoschus esculentus* L. Moench) at Humera, Northern Ethiopia. *Journal of Biology, Agriculture and Healthcare*, 6(3), 92-108.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2025): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/129134>